

Ultrasound of Common Elbow Pathology

Jon A. Jacobson, MD
FACR, FSRU, FAIM, RMSK

Musculoskeletal Radiologist
Lenox Hill Radiology, NYC

1

Disclosures

- Consultant: Bioclinica
- Contractor: POCUS PRO
- Advisory Board: Philips
- Book Royalties: Elsevier
- Not relevant to this lecture

*Note: all images from the textbook
Fundamentals of Musculoskeletal Ultrasound
are copyrighted by Elsevier Inc.*

See www.jacobsonmskus.com for syllabus other educational material

2

Pathology:


- Joint effusion and bursa
- Tendon abnormalities
- Ligament abnormalities
- Nerve abnormalities
- Soft tissue masses

3

Joint Effusion


- Olecranon recess
- Displaced hyperechoic fat pad by anechoic / hypoechoic fluid
- Best place to look with US*
- More sensitive than radiographs*

De Maeseneer, Invest Radiol
1998; 33:117



4


Joint Effusion: anterior elbow




5

Complicated Fluid vs. Synovitis


- Both may appear hypo- or isoechoic
- *Findings that suggest effusion:*
- Displacement with transducer pressure
- Joint recess collapse w/ joint movement
- Negative flow on color Doppler imaging
- Swirling with transducer pressure




6




7



8




9




10




11




12




13




14



15



16



17


Biceps Brachii Tears:

- Diagnosis of full-thickness tear versus partial-thickness tear:
 - 95% sensitivity
 - 71% specificity
 - 91% accuracy
- Shadowing: important indirect sign of tendon retraction

da Gama Lobo et al., Am J Roentgenol 2013; 200:158

18


Biceps Brachii Tendon: partial tear (short head)



Longitudinal:
 Retracted superficial short head (yellow arrows)
 Hypoechoic but intact deep long head (white arrows)

19

Biceps Tendon Tears: dynamic imaging





Partial Tear

Complete Tear

20

Bicipitoradial Bursa


- Surrounds distal biceps
 - Does not communicate to elbow joint
 - No distal biceps tendon sheath
- If distended:
 - Mechanical, inflammatory
 - Characteristic "U" shape
 - Average: 1.8 – 2.5 cm in size
 - May displace deep branch of radial nerve



Skaf AY, Radiology 1999; 212:111

21

Bicipitoradial Bursitis




22

Triceps Tear:

- Muscle injury: contusion
 - Mixed echogenicity hemorrhage
- Distal tendon injury
 - Usually partial-thickness tear
 - Superficial aspect of tendon
 - Avulsion fracture of olecranon


23

Hematoma: triceps




Longitudinal


24




25




26




27



28



29



30

Lateral Collateral Ligament Complex

- Radial collateral ligament (arrows)
- Common extensor tendon (E)
- Annular ligament (arrowhead)
- Lateral ulnar collateral ligament (curved arrow)

Jacobson J. et al. J Ultrasound Medicine 2013; 33:1041

31

Lateral Collateral Ligament Complex

- Common extensor tendon (curved arrows)
- Radial collateral ligament (arrowheads)
- Annular ligament (a)

Jacobson J. et al. J Ultrasound Medicine 2014; 33:1041

Common Extensor Tendon Removed

32

Common Extensor Tendon: tendinosis

Note: normal radial collateral ligament (white arrow)

33

Common Extensor Tendon

Patient #1
Tendinosis

Patient #2
Interstitial Tear

34

Pathology:

- Joint effusion and bursa
- Tendon abnormalities
- Ligament abnormalities
- Nerve abnormalities
- Soft tissue masses

35

Ulnar Collateral Ligament Tear


Long Axis

Normal

T1w Coronal post-gadolinium

T2w Coronal post-gadolinium

36



37

Ulnar Collateral Ligament

- Valgus stress: 30 degrees elbow flexion
 - Unlock the olecranon
 - Stress: UCL anterior bundle
- Gravity stress is adequate, equal to Telos¹
- Ultrasound measurements:
 - Reliable and precise²

Ulnar Collateral Ligament: partial tear

¹Harada M et al. J Sho Elb Surg 2014; 23:561
²Bica D et al. J Ultrasound Med 2015; 34:371

38

Ulnar Collateral Ligament: valgus stress

- >1 mm asymmetric gapping = 87% accuracy in diagnosis of UCL tear
 - MR arthrography accuracy = 88%
 - US + MR arthrography: accuracy = 98%
- Asymmetric joint space widening with stress:
 - Normal: 1.3 mm or less
 - Partial tear: 1.2 – 3.0 mm
 - Full thickness tear: 2.8 – 4.8 mm

Roedl JB et al. Radiology 2016; 279:827.

39

Ulnar Collateral Ligament: laxity

With valgus stress

40

Pathology:

- Joint effusion and bursa
- Tendon abnormalities
- Ligament abnormalities
- **Nerve abnormalities**
- Soft tissue masses

41

Ulnar Nerve: anatomy

- Behind medial epicondyle of humerus:
 - Cubital tunnel retinaculum or Osborne fascia
- Distal to epicondyle:
 - True cubital tunnel
 - Between ulnar and humeral heads: flexor carpi ulnaris
 - Under arcuate ligament

Martinoli, C. et al. Radiographics 2000;20:S199-S217

42

Ulnar Nerve: cubital tunnel syndrome

- Hypoechoic and enlarged
 - > 9 mm² area¹
 - 2.8x area compared to proximal²
- Mild hypoechoogenicity alone: may be normal
- Causes:
 - Idiopathic, overuse, joint process
 - Anconeus epitrochlearis: compression
 - Normal variant accessory muscle

¹Thoirs K et al. J Ultrasound Med 2008; 27:737
²Yoon JS et al. Muscle Nerve 2008; 38:1231

43

Anconeus Epitrochlearis

- Normal variant: 34% of population
- Roof of cubital tunnel:
 - Residual muscle
 - In absence of normal attrition forming Osborn fascia
- Secondary ulnar nerve entrapment
- Diagnose in elbow extension!

Sem Musculoskel Radiol 2000; 14:814:473

44

Isolated Ulnar Nerve Dislocation

Okamoto, J Hand Surg 2000; 25B:85

*Asymptomatic finding in 20%

45

Snapping Triceps Syndrome

- Ulnar nerve and medial triceps dislocate over apex of medial epicondyle
- Ulnar nerve and medial triceps remain in contact with each other
- Palpable snap felt through transducer

Jacobson JA et al. Radiology 2001; 220:601

46

Radial Nerve: deep branch

- Supinator syndrome:
 - Motor deficits (wrist, finger extension)
 - Abnormal electrodiagnostic studies
 - Nerve enlargement: entrapment
- Radial tunnel syndrome:
 - Pain, no motor deficits, normal EMG
 - Muscle denervation on MRI
 - No nerve enlargement

Ferdinand BD et al. Radiology 2006; 240:161

47

Supinator Syndrome: deep br. radial nv.

Abnormal Normal

48



Pathology:

- Joint effusion and bursa
- Tendon abnormalities
- Ligament abnormalities
- Nerve abnormalities
- **Soft tissue masses**

49

Cat scratch disease

- Animal scratch: usually a cat
 - *Bartonella henselae*
- Child or adolescent:
 - Most common
- Elbow:
 - Lymphadenopathy
 - Epitrochlear lymph node (medial)



50

Take-home Points:

- Joint: aspirate if concern for infection
- Biceps and triceps:
 - Anatomy explains partial-thickness tears
- Nerves: don't forget to look
- Dynamic imaging
 - Ulnar nerve dislocation, snapping triceps
 - Ulnar collateral ligament evaluation

51

Thank you!

Syllabus on line and other educational material:
www.jacobsonmksus.com

Twitter handle: @jjacobsn

52